

Research Article

# Meiotic Chromosomes Behaviour in *Chlorophytum stenopetalum* Bak

Omokanye SB<sup>1\*</sup>, Mustapha OT<sup>2</sup>, Kolawole SO<sup>3</sup>

<sup>1</sup>Department of Plant Biology, Faculty of Science, University of Ilorin, National Open University of Nigeria Offa Community Study Center Offa, Kwara State, Nigeria.

<sup>2</sup>Department of Plant Biology, Faculty of Science, University of Ilorin, Ilorin, Nigeria.

<sup>3</sup>Department of Plant Biology, Faculty of Science, Federal University Kashere, Nigeria

**\*Address for Correspondence:** Omokanye Babaunde Sikiru, Department of Plant Biology, National Open University of Nigeria, Offa community Study Center, Offa Kwara State, Nigeria; E mail: omokanyebabatundesikiru@yahoo.com and omokanyesikiru072@gmail.com

**Received:** 10 September 2020; **Accepted:** 15 October 2020; **Published:** 16 October 2020

**Citation of this article:** Omokanye SB, Mustapha OT, Kolawole SO (2020) Meiotic Chromosomes Behaviour in *Chlorophytum stenopetalum* Bak. Rea Int J of Plant Sci and Ecol. 1(1): 001-003. DOI: 10.37179/rijpse.000001.

**Copyright:** © 2020 Omokanye SB, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

Chromosomes behavior at meiosis in *Chlorophytum stenopetalum* Bak. was investigated especially at pachytene /diakinesis and metaphase. The flower buds at right age were harvested, fixed in Cornoy's solution (3 part of absolute alcohol and 1 part of acetic alcohol) and preserved in a refrigerator at -4 for at least thirty minutes. The flower buds were then hydrolyzed in 10% HCl for 3-5 minutes. Prepared slides were viewed using an Armscope microscope equipped with digital automatic camera. At diakinesis seven bivalents (7 II) were predominantly observed (74.2%). Chromosomal stickiness is quite evidence. In addition, cross configuration and its resultant, ring formation at metaphase indicate the presence of translocation heterozygosity in the chromosomes of the species investigated. These abnormalities are likely to affect microsporogenesis and pollen viability.

**Keywords:** Translocation heterozygosity, Pachytene, Diakinesis, *Chlorophytum stenopetalum* meiosis.

## Introduction

*Chlorophytum* Ker-Gawl. is a genus in the family Asperagaceae, sub-family Agavoideae of monocotyledonous perennial flowering plant, order Asparagales [1]. This position stated in Angiosperm Phylogenic Group III [2], 2009 is maintained the new classification [3]. The genus comprises about 234 species distributed in tropical and subtropical regions [4]. It is probable centre of origin and diversification lie in the tropical and sub-tropical Africa and Asia, where 85% of the species are found [5]. Many species of *Chlorophytum* are found on the forest floor in West Africa while several others occur in the guinea savanna regions [6] Meerts and Bjora, 2012). *Chlorophytum stenopetalum* is one of species of *Chlorophytum* found

in the guinea savanna region in Nigeria. The species is a tuberous herb, with rosette of leaves close to the ground. Leaves are linear-lanceolate and leaf margin is wavy. Inflorescence is about one fifth if leaf, bearing congested fruit. One to three inflorescences grow simultaneously from the centre of the leaves rosette, with the prime inflorescence at the middle. Roots are long bearing tubers at the median position.

The genus *Chlorophytum* is characterized by polyploid evolution having (x=7 or 8) chromosomes [6, 7]. Previous cytological works on species of *Chlorophytum stenopetalum* indicate a diploid chromosome number of 2n=14 [8-10]. The present work is aimed at verifying the existing diploid number and further elucidates chromosomes behavior by investigating meiotic process in the taxa.

## Materials and Methods

Samples of species of *Chlorophytum stenopetalum* used for these studies were collected at the onset of the rain in the month of March 2013, from a village called Shika dam, about 35km south of Ahmadu Bello University, Zaria. These materials were raised in pots containing garden soil, in the Department of Plant Biology of University of Ilorin. Most of the plants produced flowers in the month of April 2018 approximately a month after vegetative growth has been established, while others produce flowers much later.


The flower buds at different stages of development were harvested and fixed immediately in freshly prepared Cornoy's solution (3 part of absolute alcohol and 1 part of acetic alcohol) and preserved in a refrigerator at -4°C for at least thirty minutes, to allow for proper fixing of the cells and the removal of mucilage from the anther. The flower buds were then hydrolyzed in 10% HCl for 3-5 minutes to allow for easy removal of the perianth segments. Anther from hydrolyzed flower bud was squashed on a clean slide on which a drop of acetic orcein stain had been placed. Prepared slides were viewed using an arm scope microscope equipped digital automatic camera.

## Results


The meiotic cell division was studied from pachytene/diakinesis to full meiotic metaphase by observing microsporocytes. A total of 863 microsporocytes were studied. In diakinesis, the pollen mother cells (PMCs) revealed  $n=7$  (bivalents) for the taxa. These results thus complement the mitotic results reported earlier.




**Plate 1:** *Chlorophytum stenopetalum* at maturity stage.



**Plate 2:** Chromosome association (pairing) in metaphase I in the *C. stenopetalum*, showing diakinesis with seven bivalents (7 II).



**Plate (3a-d):** Chromosome association (pairing) in metaphase I in the taxon. a) Metaphase with 1IV+5II. b) Metaphase showing 2IV+3II c) Metaphase with 1VI+4II. d) Metaphase with 3IV+1II.



**Plate 4:** Meiotic irregularities: a, Chromosome pairing at pachytene stage b, first meiotic metaphase with spindle apparatus and chromosome clumping c, Cross shaped configuration at pachytene, di, Open ring configuration at meiotic metaphase ii, Twisted ring configuration at meiotic metaphase

Apart from the bivalents, the taxon presented different chromosomal associations, with varying frequencies of quadrivalent and hexavalent as shown in each of the (Plates 3) (Table 1) shows frequency of occurrence of the pairing patterns in *C. stenopetalum*.

Other meiotic irregularities observed include chromosome clumping on the spindle fibers, cross shaped configuration at the pachytene stage of meiosis and twisted ring and open ring configuration at meiotic metaphase.

**Table 1:** Chromosome pairing configurations (%) at diakinesis in *Chlorophytum stenopetalum* (I = univalents; II = bivalents; IV = tetravalents).

Taxon	Chromosome number	Pairing		Configurations (%)		
		7II	5II+1IV	4II+1VI	3II+2IV	1II+3IV
	2n=14					
<i>C.stenopetalum</i>		74.2	12.8	8.3	3.3	1.4

**Table 2:** Chromosome behavior at pachytene/diakinesis and metaphase I in *Chlorophytum stenopetalum*. Percentage (%) of observed configurations.

Taxon Observed formations at each of the stages				
	Cross configuration at pachytene	% of PMC with twisted ring	% of PMC with open ring	% Chromosome clumping
<i>C.stenopetalum</i>	25.3	12.1	21.3	40.4

## Discussion

The present study constitutes the first report on meiotic chromosome abnormalities and behavior in *Chlorophytum*

*stenopetalum* which revealed a high incidence of translocation heterozygosis with a remarkable variety of meiotic chromosome behavior of a wild population of the genus *Chlorophytum* in Nigeria. During the meiotic analysis seven bivalents were observed at diakinesis confirming that the species exhibits  $2n=14$  chromosomes. This number ( $2n=14$ ) is in accordance with the previous reports for the species (Adeyemi, 1974; Ngwa, 1979 and Omokanye, 2015). The present work thus indicates *C.stenopetalum* belongs to  $x=7$  series of the genus.

Though most of the pollen mother cells at diakinesis had regular meiotic behavior showing seven bivalents (74.2%), some of the homologous chromosomes paired up forming multivalent. In most cases, chromosomes intermesh losing their individual structure, forming a compact mass on the spindle fiber (Plate 4b), an abnormally generally referred to as chromosomal stickiness. Chromosome stickiness constitutes the highest frequency (40.4%) of abnormality observed at metaphase I. According to Pagliarini *et al.*(1993) and Muniyamma and Kameshwari (1996) the meiotic abnormalities such as tetravalents, precocious movement of chromosome, lagging chromosomes and chromatin bridges are found frequently in most of the species of *Chlorophytum*.

Chromosome stickiness is caused due to genetic and environmental factors and several agents have been reported to cause chromosome stickiness [11, 12], reported that stickiness may result from defective functioning of one or more types of specific non histone protein involved in chromosome organization, which are needed for chromosome separation and segregation.

The altered functioning of these proteins is caused by mutation in the structural genes coding for them (hereditary stickiness) or by the direct action of mutagens (induced stickiness) [13]. however, attributed meiotic irregularity such as multivalents, univalents laggard and non-disjunction in some Nigerian plants to structural heterozygosity or Polyploidization.

Evidence of translocation heterozygosis are easily detectable as cross configurations which are observed at the pachytene stage (plate 4c). This can result from segmental translocation. At metaphase, the translocation heterozygotes have problem in pairing up hence the formation of open ring configuration at meiotic metaphase (Plate 4di) and twisted ring configuration at meiotic metaphase (Plate 4dii). However the percentage of cross configuration observed (25.3%) is outstretched by the sum of total of twisted and open ring formation (33.4%), the presence of the two configurations in reasonable number establish the presence of translocation heterozygosity in the species. The spartial orientation of meiotic chromosomes of translocation heterozygotes in relation to the poles of the cell division influences the proper distribution of chromosome complements into daughter nuclei during meiosis I, and subsequent viability of spore formed by meiosis [14-19].

Meiotic irregularities in *Chlorophytum stenopetalum* is thus established (at least at metaphase I), further investigations of microsporocyte of this species is expected to elucidate effects of these anomalies on microspores formation as it affects pollen viability. These will be the focus point of next publication in this series.

## References

1. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society*, 181: 1-20. Link: <https://bit.ly/2He2cU2>
2. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* 161: 105-121. Link: <https://bit.ly/351WhcR>
3. Stevens PF (2001 onwards) Angiosperm Phylogeny Website. Version 14. Link: <https://bit.ly/3dE6dwT>
4. Gudadhe SP, Nathar VN, Dhoran VS (2012) Meiotic Abnormalities in *Chlorophytum comosum* (Thunb) Jacq. International journal of research in plant science, Universal research publication 2: 29-34. Link: <https://bit.ly/3k4hNnn>
5. Bordia PC, Joshi A, Simlot MM (1995) Safed musli In Medicinal and Aromatic Plants, (Eds, K.L. Chadha and R. Gupta). Advances in Horticulture, Malhotra Publishing House. New Delhi 429-451
6. Baldwin JT, Speece BM (1951) Cytogeography of *Chlorophytum* in Liberia. *American Journal of Botany* 38: 153-156. Link: <https://bit.ly/2H5hquz>
7. Darlington CD, Wylie AP (1955) Chromosome Atlas (2nd Ed.). George Allen and Unwin Ltd. London. Link: <https://amzn.to/2FCkygY>
8. Sharma AK, Chatterji AK (1958) Chromosome studies as a means of detecting the method of speciation in some members of Liliaceae. *Genet. Iber* 10: 149-178.
9. Sharma AK, Raju DT (1967) Cytological analysis of six species of *Chlorophytum*. *Bull. Bot. Soc Bengal* 21: 37-46.
10. Adeyemi FA (1981) Biosystematic studies in some Nigerian species of *Anthericum* Linn. and *Chlorophytum* Ker-Gawl (Unpublished Ph.D. Thesis) University of Ibadan. Ibadan. Link: <https://bit.ly/352mVSR>
11. Ngwa FC (1979) Cytogenetic studies in Nine taxa of the genus *Chlorophytum* Ker-Gawl. in Nigeria (Unpublished Ph.D. Thesis), University of Ibadan, Ibadan. Link: <https://bit.ly/2HaxrPP>
12. Omokanye SB (2015) Biosystematic studies of the genus *Chlorophytum* Ker- gawl. in the rain forest and savanna vegetation zones of Nigeria (Unpublished Ph.D. Thesis) University of Ilorin, Ilorin.
13. Pagliarini MS (2000) Meiotic behavior of economically important plant species: the relationship between fertility and male sterility. *Genet. Mol. Biol.* 23: 997-1002. Link: <https://bit.ly/2SWWFnp>
14. Gaulden ME (1975) Hypothesis: some mutagenes directly alterspecific chromosomal proteins (DNA) topoisomerase II and peripheral proteins) to produce chromosome stickiness, which causes chromosome aberrations. *Mutagenesis* 2: 337-365. Link: <https://bit.ly/2SWByS4>
15. Olorode O (1972) Cytogenetical studies on some Nigerian Androgoneae. *Nigeria J. Sci.* 6: 13-19. Link: <https://bit.ly/37cQ3JK>
16. Oyewole SO (1987) Cytotaxonomic studies in the genus *Urginea* Stein. In West Africa. III. The case of *Urginea indica* (Roxb.) Kunth in Nigeria. *Annals of the Missouri Botanical Garden* 74: 131-136. Link: <https://bit.ly/353tVyl>
17. Muniyamma M, Shiva Kameshwari MN (1996) Cytological studies in *Chlorophytum elatum*. (Liliaceae): *Mysore Univ* 34: 1-4.
18. Pagliarini MS (2000) Meiotic behavior of economically important plant species: the relationship between fertility and male sterility. *Genet. Mol. Biol.* 23: 997-1002. Link: <https://bit.ly/2SWWFnp>
19. Poulsen AD, Nordal I (2005) A phenetic analysis and revision of Guineo-Congolian rain forest taxa of *Chlorophytum* (Anthericaceae). *Botanical Journal of the Linnean Society* 148: 1-20. Link: <https://bit.ly/2H7k6rx>